



A RESEARCH REVIEW™
EDUCATIONAL SERIES



Making Education Easy

2019

About the Expert



Anand Prakash Dubey
MBBS, M.D. (Pediatrics), FIAP
Professor of Pediatrics

Dr A. P. Dubey is a Professor of Pediatrics at ESI Postgraduate Institute of Medical Sciences & Research, New Delhi. His career in Pediatrics spans over 35 years. He has recently retired as Head of the Department of Pediatrics at Maulana Azad Medical College and the associated Lok Nayak Hospital, New Delhi. He has been a thesis supervisor and examiner in Pediatric Examinations at various universities. He served as Honorary Lecturer in Pediatrics at University of Sheffield, U.K. He was awarded "Commonwealth Medical Fellowship" in Pediatric Hemato-oncology at the Children's Hospital, Sheffield (U.K) and also "Trainee Fellowship of the Indian Academy of Pediatrics". For his outstanding services, in 2010 the Delhi Govt. awarded him the "State Doctors Award". He has been Executive Editor of Indian Pediatrics (official medical journal of IAP), Convener and Chairperson of the immunization committee, Secretary and Chairperson of the subspecialty chapter of Nutrition. He has over 90 publications in national and international journals and many chapters in textbooks. He has attended many national and international conferences and presented papers.

Abbreviations used in this review:

IV = intravenous
ORS = oral rehydration salts
ORT = oral rehydration therapy
RCT = randomised controlled trial
UNICEF = United Nations International Children's Emergency Fund
WHO = World Health Organization

ABOUT RESEARCH REVIEW

Research Review is an independent medical publishing organisation producing electronic publications in a wide variety of specialist areas. Research Review scans 10,000 global medical journals to bring the most important clinical papers and advancements to your email inbox with advice and commentary from local specialists.

Publications are free to receive for Indian health care professionals, keeping them up to date with their chosen clinical area.

Research Review receives funding from a variety of sources including Government departments, pharmaceutical companies, insurers and other organisations with an interest in health. Content is created independently of sponsor companies with assistance from leading local specialists.

Educational Series are a summary of the most important international and local literature which impacts on treatment of a specific medical condition. These Reviews provide information on a disease, current treatment and local/international guidelines. They are intended as an educational tool.

Please send us any feedback or comments to

admin@researchreview.com.

Oral Rehydration Therapy for Diarrhea-Related Dehydration in India

This review discusses the oral treatment of dehydration associated with acute diarrhea in India and reports on the current World Health Organization (WHO) and United Nations International Children's Emergency Fund (UNICEF) recommendations for such therapy. Despite this effective, simple and affordable treatment for reducing diarrhea-related dehydration, it remains a leading cause of death in India in children under 5 years of age.¹⁻³ It is imperative that both healthcare providers and caregivers understand the importance of timely and appropriate oral rehydration therapy (ORT) in this group of patients.²

Introduction

Diarrhea-induced dehydration is a leading cause of morbidity and mortality in the developing world and is especially problematic in South-East Asian countries, including India.¹⁻⁴ Of the deaths occurring in children <5 years of age in 2016 as a result of diarrhea, 89.37% occurred in South Asia and sub-Saharan Africa.⁴ According to WHO data published in 2011, diarrhea was responsible for 18% of childhood deaths in low-income countries (**Figure 1**).⁵ In children, each episode of diarrhea reduces nutrition required for growth, resulting in undernutrition and a diarrhea-undernutrition cycle (**Figure 2**) that promotes further infection and increased morbidity and mortality, particularly in young children.⁶




Figure 1. Causes of child deaths in low-income countries.⁵



Figure 2. The cycle between Diarrhea and undernutrition.⁶

Management of diarrhea by replenishing the lost body fluid through oral rehydration therapy (ORT) is a simple, affordable, and effective intervention that significantly reduces diarrhea-related mortality and morbidity.¹ In the 1970s, WHO developed a simple, inexpensive iso-osmotic (311 mOsm/L) formulation of glucose-based oral rehydration salts (ORS) designed to correct dehydration and metabolic acidosis in the setting of acute diarrhea.^{3,7} Since the introduction of ORS, the global annual mortality rate for children suffering acute diarrhea and dehydration has dropped from 5 million to 0.5 million, and ORS remains the cornerstone of therapy for dehydration secondary to acute infectious diarrhea.^{3,4,7} Despite the ready availability of ORS, mortality rates in India still remain relatively high, with 102,813 children <5 years of age reported to have died from diarrhea in 2016, a mortality rate of 4.1 per 1000 live births.⁸



ORS solutions

ORS solutions are designed to contain the appropriate amount of sodium, glucose and other electrolytes, with appropriate osmolality to maximise water absorption from the gut using the principle of glucose-facilitated sodium transport (glucose enhances sodium and secondary water transport across the mucosa of the upper intestine).⁹ Optimal absorption of fluid from the small intestine is critically dependent on the composition of the rehydration solution. Fluid absorption depends on three factors: sodium concentration, glucose concentration and the luminal fluid osmolality.

WHO ORS formulation

In 1975, WHO and UNICEF agreed to promote a single solution (WHO-ORS) containing: sodium 90 mmol/L; potassium 20 mmol/L; chloride 80 mmol/L; base 30 mmol/L (bicarbonate); and 2% glucose (111 mmol/L). This solution struck a compromise between the ideal solutions for different disorders and the goal of a single formulation to simplify delivery and logistics for global use in cholera and non-cholera diarrheas.¹⁰

Alternative ORS formulations

Alternative formulations have been investigated in order to develop an ORS formulation that would decrease stool output or have other clinical benefits, with concerns that the original ORS sodium concentration was too high (90 mmol/L) and was occasionally associated with hypernatremia.¹¹

In 2002, WHO promoted a new low-sodium, low-glucose ORS formulation with an osmolality of 245 mOsm/L, which was associated with reduced need for unscheduled IV therapy, decreased stool output and less vomiting when compared with the original formulation.¹¹ This initiative was based on the findings of a number of studies including a meta-analysis of nine RCTs in children with acute diarrhea showing significantly fewer unscheduled IV infusions (OR 0.61; 95% CI 0.47-0.81), decreased stool output and less frequent vomiting than in children receiving the original WHO ORS formulation (311 mOsm/L).^{3,12} The low-osmolality formulation (**Table 1**) remains the currently recommended ORS for acute diarrhea.³

Other organisations have also promoted lower osmolality ORS formulations including the European Society of Paediatric Gastroenterology and Nutrition, which recommended an ORS containing 60 mmol/L of sodium with an osmolality between 200 and 250 mOsm/L for children in developed countries who are not malnourished.¹³ However, as the WHO 2002 Consensus meeting emphasised, while the efficacy of glucose-based ORS in children with acute non-cholera diarrhea was improved by reducing sodium to 60-75 mmol/L, glucose to 75-90 mmol/L, and total osmolality to 215 to 260 mOsm/L there was insufficient evidence to differentiate between ORS solutions containing less than 75 mmol/L of sodium versus >75 mmol/L, and there are considerable programmatic and logistic advantages inherent in using a single solution around the globe for diarrhea cases at all ages.³

While this single ORS low-osmolality formulation is recommended, WHO and UNICEF acknowledge that there is a range of criteria for acceptable ORS formulations.^{3,11} These criteria are as follows:

- The total substance concentration (including that contributed by glucose) should be within the range of 200-310 (mOsm/L)
- The individual substance concentration:
 - Glucose — should at least equal that of sodium but should not exceed 111 mmol/L
 - Sodium — should be within the range of 60-90 mmol/L
 - Potassium — should be within the range of 15-25 mmol/L
 - Citrate — should be within the range of 8-12 mmol/L
 - Chloride — should be within the range of 50-80 mmol/L

Composition of rehydration solutions

Compositions of available ORS solutions and other fluids differ with regard to concentrations of electrolytes and glucose, and total osmolality. The composition of various ORSs and other fluids are shown in **Table 1**. Fruit juices and soft drinks contain minimal sodium and have excessive glucose that results in excessive osmolality, which may worsen diarrhea.⁹ Sports drinks have varying sodium and carbohydrate levels, and are therefore inappropriate as rehydration solutions.⁹

Table 1. Composition of available ORS solutions and other fluids used for rehydration¹³⁻¹⁸

| Product | Carbohydrate g/L (mmol/L) | Sodium mmol/L | Potassium mmol/L | Chloride mmol/L | Base* mmol/L | Osmolarity mOsm/L |
|---------------------------------------------------------|-------------------------------------------------------|--------------------|--------------------|--------------------|------------------------------|----------------------|
| WHO (acceptable range) | At least equal to sodium, not greater than 111 mmol/L | Range 60-90 mmol/L | Range 15-25 mmol/L | Range 50-80 mmol/L | Range 8-12 mmol/L | Range 200-310 mOsm/L |
| WHO (original) | 20 (111) | 90 | 20 | 80 | 10 citrate 30 bicarbonate | 311 |
| WHO (low-osmolality) | 13.5 (75) | 75 | 20 | 65 | 10 | 245 |
| ESPGHAN European formula | 16 (74-111) | 60 | 20 | 60 | 10 | 240 |
| Pedialyte® | 25 (139) | 45 | 20 | 35 | 10 | 250 |
| Gastrolyte® | 17.8 (90) | 60 | 20 | 60 | 10 | 240 |
| ORSL™ Rehydrate Apple | 15 (65) | 60 | 20 | 51 | 10 | 259 |
| ORSL™ Rehydrate Orange | 15 (69) | 60 | 20 | 51 | 10 | 244 |
| Electral® | 13.5 (75) | 75 | 20 | 65 | 10 | 245 |
| Beverages not appropriate for diarrhea treatment | | | | | | |
| Coca Cola classic**® | 112 (622) | 1.6 | N/A | N/A | 13.4 | 650 |
| Gatorade® | 58.3 (322) | 20 | 3.2 | 11 | N/A | 350 |
| Apple juice | 120 (666) | 0.4 | 44 | 45 | N/A | 730 |

* actual or potential bicarbonate (e.g. citrate, lactate, phosphate, acetate)

** figures do not include electrolytes that might be present in local water used for bottling

Adapted from King (MMWR) 2003 and Sandhu 2001 and also includes data from commercial package inserts and labels.

SUBSCRIBE FREE OF CHARGE AT RESEARCH REVIEW www.researchreview.com/India

Privacy Policy: Research Review will record your email details on a secure database and will not release them to anyone without your prior approval.

Research Review and you have the right to inspect, update or delete your details at any time.

Research Review publications are intended for Indian healthcare professionals.



The role of zinc

Children with diarrhea receiving daily zinc supplementation have been shown to experience a significantly faster recovery from diarrhea and an 18% to 59% reduction in total stool output.¹⁶ The Indian Academy of Pediatrics, WHO and UNICEF endorse the use of zinc supplementation for 14 days during treatment with ORS for diarrhea.^{19,20}

Pre-mixed ORS solutions

In an attempt to improve the uptake of ORS, a number of manufacturers have developed pre-mixed ORS solutions, which meet WHO criteria and have the advantage of convenience, accurate dosing and improved taste. Pre-made liquid ORS solutions are less time consuming to prepare and have the advantage over sachets of not requiring interpretation of instructions for their preparation, ensuring no opportunity for errors in preparation and subsequent final osmolarity.¹⁴ Other major advantages of such commercially prepared ORS solutions are their availability in pre-flavoured formats using sweeteners that do not significantly alter their osmolarity, and the fact that they do not require access to clean drinking water by the caregiver.^{21,22}

Administration of ORS

ORS are produced in three dosage forms: powder, tablet and liquid.¹⁶ Powder sachets of ORS are usually made up in clean or boiled drinking water. Pre-made liquid ORS solutions are an effective, convenient and safe option, especially in situations where clean drinking water is unavailable, and caregivers often prefer such formulations.² ORS may be formulated with fruit juice to improve the taste, but the final product should be within the recommended osmolarity range (200-310 (mOsm/L)).²³

Assessment and treatment of diarrhea-related dehydration in children

The majority of childhood cases of mild-to-moderate diarrhea-related dehydration can be successfully managed with ORT via mouth or nasogastric tube, and this first-line treatment has been shown to be as effective, more straightforward, and less costly than IV rehydration.^{1,9,24} Contraindications to ORT include shock or suspected acute abdomen, but vomiting is not a contraindication.

The flow chart in **Figure 3** outlines the WHO assessment and treatment (Plans A, B and C) of diarrhea-related dehydration in children.²⁵




Figure 3. Assessment and treatment options for children with diarrhea-associated dehydration.²⁵

Plan A (for when 'no dehydration' is present) involves giving as much extra fluid as the child will take.²⁵ It is especially important to give ORS to use at home when the child has been treated with Plan B or Plan C during their healthcare visit, and when the child cannot return to a clinic if the diarrhea gets worse.²⁵ The amount of fluid administered should equal 50-100 mL after each loose stool in children up to 2 years of age, and 100-200 mL in those ≥2 years of age. This should be administered via frequent small sips and should be continued until the diarrhea stops.

Plan B (for when 'some dehydration' is present) involves the administration of recommended amounts of ORS over a 4-hour period as shown in **Table 2**.²⁵ If the child desires more ORS than is suggested, give more. This should be administered via frequent small sips and should be continued for 4 hours. After 4 hours the child should be reassessed and classified for dehydration (**Figure 3**) and the appropriate plan selected.

Table 2. Approximate amount of ORS solution to give in the first 4 hours in children with some dehydration²⁵

| Age ¹ | Up to 4 months | 4 -12 months | 12 -24 months | 2 - 5 years |
|------------------|----------------|--------------|---------------|-------------|
| Weight | <4 kg | 6 to <10 kg | 10 to <12 kg | 12 to <20kg |
| Fluid in ML | 200-450 | 450-800 | 800-960 | 960-1600 |

¹ Use the patient's age only when the weight is unknown. The approximate amount of ORS required (in mL) can also be calculated by multiplying the patient's weight (in kg) times 75

Plan C (for when 'severe dehydration' is present) involves administering IV fluids immediately if possible and having the child consume ORS while the drip is set up if they are able to drink (otherwise when they are able to drink, administer ORS at a rate of 5 mL/kg/hour). If it is not possible to administer IV fluids, ORS may be administered via a nasogastric tube (or by mouth) at a rate of 20 mL/kg/hour for 6 hours (total 120 mL/kg). After 6 hours, the child should be reassessed and classified for dehydration (**Figure 3**) and the appropriate plan selected. Urgent referral to hospital for IV therapy may be required in severe cases.

For ongoing care, emphasis should be placed on educating caregivers on appropriate rehydration in diarrheal diseases explaining that administration of water, or carbohydrate-only containing drinks, does not promote adequate fluid absorption in dehydrated individuals and that drinks with a high glucose content and unbalanced carbohydrate/sodium ratios are not suitable for rehydration as they may worsen diarrhea through osmotic mechanisms.²⁶

Issues for rehydration in India

In 2004 an Indian National Expert Group created by the Indian Ministry of Health recommended that a single universal ORS solution containing sodium 75 mmol/L and glucose 75 mmol/L, with an osmolarity of 245 mOsm/L was acceptable for all ages and all types of diarrhea.²⁰ In 2012, a survey revealed that the prescription of ORS reported by caregivers in India is low, ranging between 19% and 47%, despite 90% of private providers perceiving ORS favourably and 89% acknowledging it as the recommended first-line treatment of diarrhea.² The commonly reported caregiver perception that diarrhea is not a life-threatening disease and the frequently employed 'wait and watch' approach, where medical attention is only sought when fever, vomiting or severe dehydration develop, is considered a major reason for the poor uptake of ORS in developing countries.² Other contributing factors include the poor palatability of some ORS solutions, cost and accessibility.²¹



A recent observational study conducted at an Indian medical college hospital catering mainly for the middle and lower class population, identified insufficient caregiver knowledge regarding adequate hand washing (51%) incorrect dilution of ORS (70%), and the use of unsafe water in the preparation of ORS (29%), as correctable factors associated with home-based ORT for the management of diarrhea.¹ Another study revealed that only one-fifth of mothers were aware of zinc supplementation and only 7% of children with diarrhea received such treatment.²⁴

There is a clear and urgent need for caregiver education regarding the importance of ORT and zinc supplementation in children with mild-to-moderate dehydration, with an emphasis on the importance of correct preparation and administration of ORS.¹

Key measures to prevent diarrhea

WHO cite the following as key measures for the prevention of diarrhea:²⁵

- Access to safe drinking water
- Use of improved sanitation
- Hand washing with soap
- Exclusive breastfeeding for the first six months of life
- Good personal and food hygiene
- Health education about how infections spread
- Rotavirus vaccination.

EXPERT'S CONCLUDING COMMENTS

This article describes Oral Rehydration Therapy (ORT) in children suffering from acute diarrhea and dehydration in a practical way. Current WHO Oral Rehydration Solutions (ORS) offer the most appropriate water and electrolyte replenishment in all types of diarrheal dehydration. ORT is the most cost effective and easily accessible intervention to decrease the morbidity and mortality in such cases. All children suffering from diarrhea related dehydration should be offered ORS as the primary therapeutic intervention under all circumstances.

TAKE-HOME MESSAGES

- Key measures including improved hygiene and sanitation, and vaccination for rotavirus should be implemented for the prevention of diarrhea
- Dehydration should be prevented and treated in children with diarrhea by administering an appropriate ORS solution
- WHO/UNICEF recommend the single low-osmolarity ORS formulation, but acknowledge that there is a range of acceptable ORS formulations
- Most cases of diarrhea-related dehydration can be successfully treated with ORS
- Zinc supplementation should be provided for a duration of 14 days in children and infants receiving ORS for diarrhea
- Pre-made liquid ORS solutions may be an effective, palatable and convenient option, especially in situations where clean drinking water is unavailable.

REFERENCES:

1. Gandra NR and Farooqui D. Diarrhoea, nutrition and oral rehydration therapy: awareness, attitude and practices among mothers of children under five years. *Int J Contemp Pediatr*. 2017;4(6):1995-2001
2. Clinton Health Access Initiative New Delhi, India. The private sector market for diarrhea treatment in India. 2012. Available from: <http://www.zinctaskforce.org/wp-content/uploads/2011/06/The-Private-Sector-Market-for-Diarrhea-Treatment-in-India.pdf> (Accessed Dec 2018)
3. World Health Organization. WHO Drug Information 2002;16(2). Available from: <http://apps.who.int/medicinedocs/pdf/s4950e/s4950e.pdf> (Accessed Dec 2018)
4. GBD 2016 Causes of Death Collaborators. Global, regional, and national age-sex specific mortality for 264 causes of death, 1980–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 2017; 390: 1151–210
5. World Health Organization. World Health Statistics 2011. Available from: <https://www.who.int/whosis/whostat/2011/en/> (Accessed Dec 2018)
6. Office of Global Health and HIV (OGHH) Office of Overseas Programming & Training Support (OPATS) Integrating WASH and Nutrition, Water, Sanitation and Hygiene Training Package. Available from: <https://slideplayer.com/slide/3903727/> (Accessed Dec 2018)
7. Binder HJ et al. Oral rehydration therapy in the second decade of the twenty-first century. *Curr Gastroenterol Rep*. 2014;16:376
8. Pneumonia & Diarrhea Progress Report 2018. <https://www.jhsph.edu/ivac/wp-content/uploads/2018/11/PDPR2018.pdf> (Accessed Dec 2018)
9. Infants and Children: Management of Acute Gastroenteritis, Fourth Edition. NSW Government Health Guidelines. 2014 Available from: https://www1.health.nsw.gov.au/pds/ActivePDSDocuments/GL2014_024.pdf (Accessed Dec 2018)
10. MMWR Recommendations and Reports 1992; 41;RR-16:001. Available from: <https://www.cdc.gov/mmwr/preview/mmwrhtml/00018677.htm> (Accessed Dec 2018)
11. World Health Organization and United Nations International Children's Emergency Fund. Oral rehydration salts. Production of the new ORS. WHO 2006. Available from: http://apps.who.int/iris/bitstream/handle/10665/69227/WHO_FCH_CAH_06.1.pdf?sequence=1 (Accessed Dec 2018)
12. Hahn S et al. Reduced osmolarity oral rehydration solution for treating dehydration due to diarrhoea in children: systematic review. *BMJ*. 2001;323(7304):81-5
13. Guarino A et al. European Society for Pediatric Gastroenterology, Hepatology, and Nutrition/European Society for Pediatric Infectious Diseases evidence-based guidelines for the management of acute gastroenteritis in children in Europe: update 2014. *J Pediatr Gastroenterol Nutr*. 2014;59(1):132-52
14. King CK et al. Managing acute gastroenteritis among children. MMWR Recommendations and reports 2003;52(RR16):1-16 Available from: <https://www.cdc.gov/mmwr/preview/mmwrhtml/rr5216a1.htm> (Accessed Dec 2018)
15. Atia AN et al. Oral rehydration solutions in non-cholera diarrhoea: A review. *Am J Gastroenterol*. 2009;104:2596-604
16. Sandhu BK. Practical guidelines for the management of gastroenteritis in children. *J Pediatr Gastroenterol Nutr* 2001;33(Suppl 2):S36--9
17. JNJ -ORSL Rehydrate chemical analysis (data on file)
18. Electral packaging <http://www.fdcindia.com/formulations.php> (Accessed Dec 2018)
19. World Health Organization. Implementing the new recommendations on the clinical management of diarrhoea. Guidelines for policy makers and programme managers. 2006. Available from: http://apps.who.int/iris/bitstream/handle/10665/43456/9241594217_eng.pdf?sequence=1 (Accessed Dec 2018)
20. Bhatnagar S et al. IAP Guidelines 2006 on Management of Acute Diarrhea. *Indian Pediatrics* 2007;44:380-389. Available from: <https://indianpediatrics.net/may2007/may-380-389.htm> (Accessed Dec 2018)
21. Freedman SB et al. Assessing the palatability of oral rehydration solutions in school-aged children. A randomised crossover trial. *Arch Pediatr Adolesc Med*. 2010;164(8):696-702
22. Santosham M and Greenough WB. Oral rehydration therapy: A global perspective. *J Pediatrics* 1991;118(4 Pt 2):S44-51
23. te Loo DM et al. The effect of flavoring oral rehydration solution on its composition and palatability. *J Pediatr Gastroenterol Nutr*. 2004 Nov;39(5):545-8
24. Hartling L et al. Oral versus intravenous rehydration for treating dehydration due to gastroenteritis in children. *Cochrane Database Syst Rev*. 2006;Jul 19;(3):CD004390
25. World Health Organization. Manual for the Health Care of Children in Humanitarian Emergencies. 2008. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK143745/> (Accessed Dec 2018)
26. Guarino A et al. Oral rehydration solution – An essential therapy for childhood gastroenteritis. *JAMA Pediatrics* 2018;172(10):991
27. Ali F et al. Assessing critical gaps in implementation of WHO and UNICEF'S 7-point diarrhea control and prevention strategy in Uttar Pradesh, India. *Tropical Med Pub Health* 2017;10(3):571-79
28. World Health Organization. Diarrhoeal disease. 2017. Available from: <https://www.who.int/news-room/fact-sheets/detail/diarrhoeal-disease> (Accessed Dec 2018)

SUBSCRIBE FREE of CHARGE AT RESEARCH REVIEW www.researchreview.com/India



This publication has been created with an educational grant from Johnson and Johnson Asia Pacific. The content is entirely independent and based on published studies and the writer and commentators' opinions.